

The Cauchy problem for higher-order linear partial differential equation

Guangqing Bi ^{a)†}, Yuekai Bi ^{b)}

^{b)} School of Electronic and Information Engineering, BUAA, Beijing 100191, China

E-mail: yuekaifly@163.com

For the linear partial differential equation $P(\partial_x, \partial_t)u = f(x, t)$, where $x \in \mathbb{R}^n$, $t \in \mathbb{R}^1$, with $P(\partial_x, \partial_t)$ is $\prod_{i=1}^m (\frac{\partial}{\partial t} - a_i P(\partial_x))$ or $\prod_{i=1}^m (\frac{\partial^2}{\partial t^2} - a_i^2 P(\partial_x))$, the authors give the analytic solution of the cauchy problem using the abstract operators $e^{tP(\partial_x)}$ and $\frac{\sinh(tP(\partial_x)^{1/2})}{P(\partial_x)^{1/2}}$. By representing the operators with integrals, explicit solutions are obtained with an integral form of a given function.

Keywords: cauchy problem, partial differential equation, abstract operators

MSC(2000): 35G10; 35G05

1. Introduction and main results

In 1997, Guangqing Bi first introduced the concept of abstract operators in reference [1], and determined the algorithms of abstract operators $e^{tP(\partial_x)}$ and $\frac{\sinh(tP(\partial_x)^{1/2})}{P(\partial_x)^{1/2}}$. Using this type of operators, in reference [2] the author has obtained the following results:

Theorem BI1. Let $a_1, a_2, \dots, a_m$ be arbitrary real or complex numbers different from each other, $P(\partial_x)$ be a partial differential operator of any order, then $\forall f(x, t) \in C^\infty(\mathbb{R}_x^n \times \mathbb{R}_t^1)$, we have

$$\begin{cases} \prod_{i=1}^m (\frac{\partial}{\partial t} - a_i P(\partial_x))u = f(x, t), & x \in \mathbb{R}^n, \quad t \in \mathbb{R}^1. \\ \frac{\partial^j u}{\partial t^j} \Big|_{t=0} = 0, & j = 0, 1, 2, \dots, m-1. \end{cases} \quad (1)$$

$$u(x, t) = \int_0^t \int_0^{t-\tau} \frac{(t - \tau - \tau')^{m-2}}{(m-2)!} \sum_{j=1}^m \frac{a_j^{m-1}}{\prod_{i=1, i \neq j}^m (a_j - a_i)} e^{\tau' a_j P(\partial_x)} f(x, \tau) d\tau' d\tau. \quad (2)$$

[†]Corresponding author. E-mail: yananbiguangqing@sohu.com

Theorem BI2. Let $a_1, a_2, \dots, a_m$ be arbitrary real or complex numbers different from each other, $P(\partial_x)$ be a partial differential operator of any order, then $\forall f(x, t) \in C^\infty(\mathbb{R}_x^n \times \mathbb{R}_t^1)$, we have

$$\begin{cases} \prod_{i=1}^m (\frac{\partial^2}{\partial t^2} - a_i^2 P(\partial_x)) u = f(x, t), & x \in \mathbb{R}^n, \quad t \in \mathbb{R}^1. \\ \frac{\partial^j u}{\partial t^j} \Big|_{t=0} = 0, & j = 0, 1, 2, \dots, 2m-1. \end{cases} \quad (3)$$

$$u(x, t) = \int_0^t \int_0^{t-\tau} \frac{(t-\tau-\tau')^{2m-3}}{(2m-3)!} \sum_{j=1}^m \frac{a_j^{2m-2}}{\prod_{i \neq j}^m (a_j^2 - a_i^2)} \frac{\sinh(\tau' a_j P(\partial_x)^{1/2})}{a_j P(\partial_x)^{1/2}} f(x, \tau) d\tau' d\tau. \quad (4)$$

In reference [3] Guangqing Bi has obtained the following results:

Theorem BI3. Let $m \in \mathbb{N}$ and $m > 1$, $P(\partial_x)$ be a partial differential equation of any order. Then $\forall f(x, t) \in C^\infty(\mathbb{R}_x^n \times \mathbb{R}_t^1)$, we have

$$\begin{cases} \left(\frac{\partial^2}{\partial t^2} - P(\partial_x) \right)^m u = f(x, t), & x \in \mathbb{R}^n, \quad t \in \mathbb{R}^1. \\ \frac{\partial^j u}{\partial t^j} \Big|_{t=0} = 0, & j = 0, 1, 2, \dots, 2m-1. \end{cases} \quad (5)$$

$$u(x, t) = \int_0^t \int_0^{t-\tau} \frac{((t-\tau)^2 - \tau'^2)^{m-2}}{(2m-2)!!(2m-4)!!} \frac{\sinh(\tau' P(\partial_x)^{1/2})}{P(\partial_x)^{1/2}} f(x, \tau) d\tau' d\tau. \quad (6)$$

By combining the abstract operators and Laplace transform, the authors have obtained the following results in reference [4]:

Theorem BI4. Let $m \in \mathbb{N}$ and $m > 1$, $P(\partial_x)$ be a partial differential equation of any order. Then $\forall f(x, t) \in C^\infty(\mathbb{R}_x^n \times \mathbb{R}_t^1)$, $\varphi_j(x) \in C^\infty(\mathbb{R}^n)$, we have

$$\begin{cases} \left(\frac{\partial^2}{\partial t^2} - P(\partial_x) \right)^m u = f(x, t), & x \in \mathbb{R}^n, \quad t \in \mathbb{R}^1. \\ \frac{\partial^j u}{\partial t^j} \Big|_{t=0} = \varphi_j(x), & j = 0, 1, 2, \dots, 2m-1. \end{cases} \quad (7)$$

$$\begin{aligned} u(x, t) &= \int_0^t \int_0^{t-\tau} \frac{((t-\tau)^2 - \tau'^2)^{m-2}}{(2m-2)!!(2m-4)!!} \frac{\sinh(\tau' P(\partial_x)^{1/2})}{P(\partial_x)^{1/2}} f(x, \tau) \tau' d\tau' d\tau \\ &+ \sum_{k=0}^{m-1} (-1)^k \binom{m}{k} P(\partial_x)^k \sum_{j=0}^{2m-1-2k} \frac{\partial^{2m-1-2k-j}}{\partial t^{2m-1-2k-j}} \int_0^t \frac{(t^2 - \tau^2)^{m-2} \tau}{(2m-2)!!(2m-4)!!} \\ &\times \frac{\sinh(\tau P(\partial_x)^{1/2})}{P(\partial_x)^{1/2}} \varphi_j(x) d\tau. \end{aligned} \quad (8)$$

Using the same method, we have obtained the following results in this paper:

Theorem 1. Let $a_1, a_2, \dots, a_m$ be arbitrary real or complex roots different from each other for $b_0 + b_1x + b_2x^2 + \dots + b_mx^m = 0$, and $P(\partial_x, \partial_t)$ be a partial differential operator defined by

$$P(\partial_x, \partial_t) = \sum_{k=0}^m b_k P(\partial_x)^{m-k} \frac{\partial^k}{\partial t^k}, \quad x \in \mathbb{R}^n, \quad t \in \mathbb{R}^1, \quad 1 < m \in \mathbb{N}.$$

Where $P(\partial_x)$ is a partial differential operator of any order. Then $\forall f(x, t) \in C^\infty(\mathbb{R}_x^n \times \mathbb{R}_t^1)$, $\varphi_r(x) \in C^\infty(\mathbb{R}^n)$, we have

$$\begin{cases} P(\partial_x, \partial_t)u = f(x, t), & x \in \mathbb{R}^n, \quad t \in \mathbb{R}^1. \\ \left. \frac{\partial^r u}{\partial t^r} \right|_{t=0} = \varphi_r(x), & r = 0, 1, 2, \dots, m-1. \end{cases} \quad (9)$$

$$\begin{aligned} u(x, t) &= \int_0^t \int_0^{t-\tau} \frac{(t-\tau-\tau')^{m-2}}{(m-2)!} \sum_{j=1}^m \frac{a_j^{m-1}}{\prod_{i \neq j}^m (a_j - a_i)} e^{\tau' a_j P(\partial_x)} f(x, \tau) d\tau' d\tau \\ &+ \sum_{k=1}^m b_k P(\partial_x)^{m-k} \sum_{r=0}^{k-1} \frac{\partial^{k-1-r}}{\partial t^{k-1-r}} \int_0^t \frac{(t-\tau)^{m-2}}{(m-2)!} \\ &\times \sum_{j=1}^m \frac{a_j^{m-1}}{\prod_{i \neq j}^m (a_j - a_i)} e^{\tau a_j P(\partial_x)} \varphi_r(x) d\tau. \end{aligned} \quad (10)$$

Theorem 2. Let $a_1, a_2, \dots, a_m$ be arbitrary real or complex roots different from each other, satisfy $\sum_{k=0}^m b_{2k} x^{2k} = \prod_{i=1}^m (x^2 - a_i^2)$, and $P(\partial_x, \partial_t)$ be a partial differential operators defined by

$$P(\partial_x, \partial_t) = \sum_{k=0}^m b_{2k} P(\partial_x)^{m-k} \frac{\partial^{2k}}{\partial t^{2k}}, \quad x \in \mathbb{R}^n, \quad t \in \mathbb{R}^1, \quad 1 < m \in \mathbb{N}.$$

Where $P(\partial_x)$ be a partial differential operator of any order, then $\forall f(x, t) \in C^\infty(\mathbb{R}_x^n \times \mathbb{R}_t^1)$, $\varphi_r(x) \in C^\infty(\mathbb{R}^n)$, we have

$$\begin{cases} P(\partial_x, \partial_t)u = f(x, t), & x \in \mathbb{R}^n, \quad t \in \mathbb{R}^1. \\ \left. \frac{\partial^r u}{\partial t^r} \right|_{t=0} = \varphi_r(x), & r = 0, 1, 2, \dots, 2m-1. \end{cases} \quad (11)$$

$$\begin{aligned} u(x, t) &= \int_0^t \int_0^{t-\tau} \frac{(t-\tau-\tau')^{2m-3}}{(2m-3)!} \sum_{j=1}^m \frac{a_j^{2m-2}}{\prod_{i \neq j}^m (a_j^2 - a_i^2)} \frac{\sinh(\tau' a_j P(\partial_x)^{1/2})}{a_j P(\partial_x)^{1/2}} f(x, \tau) d\tau' d\tau \\ &+ \sum_{k=1}^m b_{2k} P(\partial_x)^{m-k} \sum_{r=0}^{2k-1} \frac{\partial^{2k-1-r}}{\partial t^{2k-1-r}} \int_0^t \frac{(t-\tau)^{2m-3}}{(2m-3)!} \\ &\times \sum_{j=1}^m \frac{a_j^{2m-2}}{\prod_{i \neq j}^m (a_j^2 - a_i^2)} \frac{\sinh(\tau a_j P(\partial_x)^{1/2})}{a_j P(\partial_x)^{1/2}} \varphi_r(x) d\tau. \end{aligned} \quad (12)$$

2. Proof of theorems

According to the Theorem BI1 and Theorem BI2, we just need to prove the following corollary of Theorem 1 and Theorem 2:

Corollary 1. Let $a_1, a_2, \dots, a_m$ be arbitrary real or complex roots different from each other for $b_0 + b_1x + b_2x^2 + \dots + b_mx^m = 0$, and $P(\partial_x, \partial_t)$ be a partial differential operators defined by

$$P(\partial_x, \partial_t) = \sum_{k=0}^m b_k P(\partial_x)^{m-k} \frac{\partial^k}{\partial t^k}, \quad x \in \mathbb{R}^n, \quad t \in \mathbb{R}^1, \quad 1 < m \in \mathbb{N}.$$

Where $P(\partial_x)$ be a partial differential operator of any order, then for $\forall \varphi_r(x) \in C^\infty(\mathbb{R}^n)$, we have

$$\begin{cases} P(\partial_x, \partial_t)u = 0, & x \in \mathbb{R}^n, \quad t \in \mathbb{R}^1. \\ \left. \frac{\partial^r u}{\partial t^r} \right|_{t=0} = \varphi_r(x), & r = 0, 1, 2, \dots, m-1. \end{cases} \quad (13)$$

$$u(x, t) = \sum_{k=1}^m b_k P(\partial_x)^{m-k} \sum_{r=0}^{k-1} \frac{\partial^{k-1-r}}{\partial t^{k-1-r}} \int_0^t \frac{(t-\tau)^{m-2}}{(m-2)!} \sum_{j=1}^m \frac{a_j^{m-1} e^{\tau a_j P(\partial_x)}}{\prod_{i=1, i \neq j}^m (a_j - a_i)} \varphi_r(x) d\tau. \quad (14)$$

Proof. Considering initial conditions, the Laplace transform of the Eq (13) with respect to t is

$$\begin{aligned} \sum_{k=0}^m b_k P(\partial_x)^{m-k} (s^k U(x, s) - \sum_{r=0}^{k-1} s^{k-1-r} \varphi_r(x)) &= 0, \\ \prod_{i=1}^m (s - a_i P(\partial_x)) U(x, s) - \sum_{k=1}^m b_k P(\partial_x)^{m-k} \sum_{r=0}^{k-1} s^{k-1-r} \varphi_r(x) &= 0. \end{aligned}$$

Where $U(x, s) = Lu(x, t)$. Let $G_m(\partial_x, t) = L^{-1}[1/\prod_{i=1}^m (s - a_i P(\partial_x))]$, by solving $U(x, s)$, we have its inverse Laplace transform:

$$\begin{aligned} u(x, t) &= L^{-1}U(x, s) = \sum_{k=1}^m b_k P(\partial_x)^{m-k} \sum_{r=0}^{k-1} L^{-1} \frac{s^{k-1-r}}{\prod_{i=1}^m (s - a_i P(\partial_x))} \varphi_r(x) \\ &= \sum_{k=1}^m b_k P(\partial_x)^{m-k} \sum_{r=0}^{k-1} \frac{\partial^{k-1-r}}{\partial t^{k-1-r}} G_m(\partial_x, t) \varphi_r(x). \end{aligned} \quad (15)$$

Now let us solve $G_m(\partial_x, t)$. Considering initial conditions, the Laplace transform of the Eq (1) with respect to t is

$$\prod_{i=1}^m (s - a_i P(\partial_x)) U(x, s) = F(x, s), \quad F(x, s) = Lf(x, t).$$

By solving $U(x, s)$ and using the convolution theorem, we have its inverse Laplace transform:

$$u(x, t) = L^{-1}U(x, s) = L^{-1} \frac{1}{\prod_{i=1}^m (s - a_i P(\partial_x))} F(x, s) = G_m(\partial_x, t) * f(x, t).$$

By comparing (2) with $u(x, t) = G_m(\partial_x, t) * f(x, t)$, we have the expression of the abstract operator $G_m(\partial_x, t)$:

$$G_m(\partial_x, t) = \int_0^t \frac{(t - \tau)^{m-2}}{(m-2)!} \sum_{j=1}^m \frac{a_j^{m-1}}{\prod_{\substack{i=1 \\ i \neq j}}^m (a_j - a_i)} e^{\tau a_j P(\partial_x)} d\tau. \quad (16)$$

Applying (16) to (15), thus the Corollary 1 is proved.

Corollary 2. Let $a_1, a_2, \dots, a_m$ be arbitrary real or complex roots different from each other, which satisfy $\sum_{k=0}^m b_{2k} x^{2k} = \prod_{i=1}^m (x^2 - a_i^2)$, and $P(\partial_x, \partial_t)$ be a partial differential operator defined by

$$P(\partial_x, \partial_t) = \sum_{k=0}^m b_{2k} P(\partial_x)^{m-k} \frac{\partial^{2k}}{\partial t^{2k}}, \quad x \in \mathbb{R}^n, \quad t \in \mathbb{R}^1, \quad 1 < m \in \mathbb{N}.$$

Where $P(\partial_x)$ is a partial differential operator of any order, then $\forall \varphi_r(x) \in C^\infty(\mathbb{R}^n)$, we have

$$\begin{cases} P(\partial_x, \partial_t)u = 0, & x \in \mathbb{R}^n, \quad t \in \mathbb{R}^1. \\ \left. \frac{\partial^r u}{\partial t^r} \right|_{t=0} = \varphi_r(x), & r = 0, 1, 2, \dots, 2m-1. \end{cases} \quad (17)$$

$$\begin{aligned} u(x, t) &= \sum_{k=1}^m b_{2k} P(\partial_x)^{m-k} \sum_{r=0}^{2k-1} \frac{\partial^{2k-1-r}}{\partial t^{2k-1-r}} \int_0^t \frac{(t - \tau)^{2m-3}}{(2m-3)!} \\ &\quad \times \sum_{j=1}^m \frac{a_j^{2m-2}}{\prod_{\substack{i=1 \\ i \neq j}}^m (a_j^2 - a_i^2)} \frac{\sinh(\tau a_j P(\partial_x)^{1/2})}{a_j P(\partial_x)^{1/2}} \varphi_r(x) d\tau. \end{aligned} \quad (18)$$

Proof. Considering initial conditions, the Laplace transform of the Eq (17) with respect to t is

$$\sum_{k=0}^m b_{2k} P(\partial_x)^{m-k} (s^{2k} U(x, s) - \sum_{r=0}^{2k-1} s^{2k-1-r} \varphi_r(x)) = 0,$$

$$\prod_{i=1}^m (s^2 - a_i^2 P(\partial_x)) U(x, s) - \sum_{k=1}^m b_{2k} P(\partial_x)^{m-k} \sum_{r=0}^{2k-1} s^{2k-1-r} \varphi_r(x) = 0.$$

Where $U(x, s) = Lu(x, t)$. Let $G_m(\partial_x, t) = L^{-1}[1/\prod_{i=1}^m(s^2 - a_i^2 P(\partial_x))]$, by solving $U(x, s)$, we have its inverse Laplace transform:

$$\begin{aligned} u(x, t) &= L^{-1}U(x, s) = \sum_{k=1}^m b_{2k} P(\partial_x)^{m-k} \sum_{r=0}^{2k-1} L^{-1} \frac{s^{2k-1-r}}{\prod_{i=1}^m (s^2 - a_i^2 P(\partial_x))} \varphi_r(x) \\ &= \sum_{k=1}^m b_{2k} P(\partial_x)^{m-k} \sum_{r=0}^{2k-1} \frac{\partial^{2k-1-r}}{\partial t^{2k-1-r}} G_m(\partial_x, t) \varphi_r(x). \end{aligned} \quad (19)$$

Now let us solve the $G_m(\partial_x, t)$. Considering initial conditions, the Laplace transform of the Eq (3) with respect to t is

$$\prod_{i=1}^m (s^2 - a_i^2 P(\partial_x)) U(x, s) = F(x, s), \quad F(x, s) = Lf(x, t).$$

By solving $U(x, s)$ and using the convolution theorem, we have its inverse Laplace transform:

$$u(x, t) = L^{-1}U(x, s) = L^{-1} \frac{1}{\prod_{i=1}^m (s^2 - a_i^2 P(\partial_x))} F(x, s) = G_m(\partial_x, t) * f(x, t).$$

By comparing (4) with $u(x, t) = G_m(\partial_x, t) * f(x, t)$, we have the expression of the abstract operator $G_m(\partial_x, t)$:

$$G_m(\partial_x, t) = \int_0^t \frac{(t - \tau)^{2m-3}}{(2m-3)!} \sum_{j=1}^m \frac{a_j^{2m-2}}{\prod_{i \neq j}^m (a_j^2 - a_i^2)} \frac{\sinh(\tau a_j P(\partial_x)^{1/2})}{a_j P(\partial_x)^{1/2}} d\tau. \quad (20)$$

Applying (20) to (19), thus the Corollary 2 is proved.

3. Examples

Theorem 3. Let $a_1, a_2, \dots, a_m$ be arbitrary real roots different from each other, which satisfy $\sum_{k=0}^m b_{2k} x^{2k} = \prod_{i=1}^m (x^2 - a_i^2)$, and $P(\partial_x, \partial_t)$ be a partial differential operator defined by

$$P(\partial_x, \partial_t) = \sum_{k=0}^m b_{2k} \Delta_n^{m-k} \frac{\partial^{2k}}{\partial t^{2k}}, \quad x \in \mathbb{R}^n, \quad t \in \mathbb{R}^1, \quad 1 < m \in \mathbb{N}.$$

Where Δ_n is an n -dimensional Laplacian, and $n - 2 = 2\nu + 1$, $\nu \in \mathbb{N}_0$, then $\forall f(x, t) \in C^\infty(\mathbb{R}_x^n \times \mathbb{R}_t^1)$, $\varphi_r(x) \in C^\infty(\mathbb{R}^n)$, we have

$$\begin{cases} P(\partial_x, \partial_t)u = f(x, t), & x \in \mathbb{R}^n, \quad t \in \mathbb{R}^1. \\ \left. \frac{\partial^r u}{\partial t^r} \right|_{t=0} = \varphi_r(x), & r = 0, 1, 2, \dots, 2m-1. \end{cases} \quad (21)$$

$$\begin{aligned}
u(x, t) = & \int_0^t d\tau \int_0^{t-\tau} d\tau' \frac{(t - \tau - \tau')^{2m-3}}{(2m-3)!} \sum_{j=1}^m \frac{a_j^{2m-2}}{\prod_{\substack{i=1 \\ i \neq j}}^m (a_j^2 - a_i^2)} \\
& \times [\underbrace{\tau' \int_0^{\tau'} \tau' d\tau' \cdots \int_0^{\tau'} \frac{(a_j^2 \Delta_n)^\nu}{S'_{n,j}}}_{\nu} \int_{S'_{n,j}} f(\xi', \tau) dS'_{n,j} \tau' d\tau' + \sum_{l=0}^{\nu-1} \frac{a_j^{2l} \tau'^{2l+1}}{(2l+1)!} \Delta_n^l f(x, \tau)] \\
& + \sum_{k=1}^m b_{2k} P(\partial_x)^{m-k} \sum_{r=0}^{2k-1} \frac{\partial^{2k-1-r}}{\partial t^{2k-1-r}} \int_0^t d\tau \frac{(t - \tau)^{2m-3}}{(2m-3)!} \sum_{j=1}^m \frac{a_j^{2m-2}}{\prod_{\substack{i=1 \\ i \neq j}}^m (a_j^2 - a_i^2)} \\
& \times [\underbrace{\tau \int_0^\tau \tau d\tau \cdots \int_0^\tau \frac{(a_j^2 \Delta_n)^\nu}{S_{n,j}}}_{\nu} \int_{S_{n,j}} \varphi_r(\xi) dS_{n,j} \tau d\tau + \sum_{l=0}^{\nu-1} \frac{a_j^{2l} \tau^{2l+1}}{(2l+1)!} \Delta_n^l \varphi_r(x)]. \quad (22)
\end{aligned}$$

Where $S'_{n,j} = 2(2\pi)^{\nu+1}(a_j \tau')^{n-1}$, $S_{n,j} = 2(2\pi)^{\nu+1}(a_j \tau)^{n-1}$, and $\xi' \in \mathbb{R}_n$ is the integral variable. The integral is on the hypersphere $(\xi'_1 - x_1)^2 + (\xi'_2 - x_2)^2 + \cdots + (\xi'_n - x_n)^2 = (a_j \tau')^2$, and $dS'_{n,j}$ is its surface element. $\xi \in \mathbb{R}_n$ is the integral variable on the hypersphere $(\xi_1 - x_1)^2 + (\xi_2 - x_2)^2 + \cdots + (\xi_n - x_n)^2 = (a_j \tau)^2$, and $dS_{n,j}$ is its surface element.

Proof. According to (30) in reference [4], we have:

$$\begin{aligned}
\frac{\sinh(t a_j \Delta_n^{1/2})}{a_j \Delta_n^{1/2}} f(x) = & \underbrace{t \int_0^t t dt \cdots \int_0^t}_{\nu} \frac{(a_j^2 \Delta_n)^\nu}{S_{n,j}} \int_{S_{n,j}} f(\xi) dS_{n,j} t dt \\
& + \sum_{l=0}^{\nu-1} \frac{a_j^{2l} t^{2l+1}}{(2l+1)!} \Delta_n^l f(x), \quad \nu = \frac{n-3}{2}. \quad (23)
\end{aligned}$$

Where Δ_n is an n -dimensional Laplacian, $n-2 = 2\nu+1$, $S_{n,j} = 2(2\pi)^{\nu+1}(a_j t)^{n-1}$. $\xi \in \mathbb{R}_n$ is the integral variable on the hypersphere $(\xi_1 - x_1)^2 + (\xi_2 - x_2)^2 + \cdots + (\xi_n - x_n)^2 = (a_j t)^2$, and $dS_{n,j}$ is its surface element.

In Theorem 2, let $P(\partial_x) = \Delta_n$, then Theorem 3 is proved by the substitution of (23).

Similarly, we can easily obtain explicit solutions of the Cauchy problem of more complex partial differential equations. For the initial-boundary value problem, the operator $P(\partial_x)$ must have the characteristic function related to boundary conditions, in order to expand the known function $f(x, \tau)$, $\varphi_r(x)$ in (10) or (12) by using the characteristic function of $P(\partial_x)$. Therefore, if the determination of the characteristic function can be ascribed to the Sturm-Liouville problem of given boundary conditions, then this initial-boundary value problem is solvable.

$P(\partial_x)$ in Theorem BI4, Theorem 1 and Theorem 2 can be variable-coefficient partial differential operators. For instance, if $P(\partial_x)$ is a self-adjoint operator defined in a Hilbert space,

then the abstract operator

$$\frac{\sinh(tP(\partial_x)^{1/2})}{P(\partial_x)^{1/2}} \quad \text{and} \quad \cosh(tP(\partial_x)^{1/2}) = \frac{\partial}{\partial t} \frac{\sinh(tP(\partial_x)^{1/2})}{P(\partial_x)^{1/2}}$$

can act on the Hilbert space, which also is a bounded operator in the Hilbert space. In this case, we can attach proper boundary conditions to the initial value problems in (7), (9) and (11). Therefore, the given function $f(x, t), \varphi_r(x)$ becomes a function with boundary conditions, and can be expressed in a Hilbert space within the given domain. In order to solve the corresponding initial-boundary value problem, we need to solve the characteristic value problem of $P(\partial_x)$ under given boundary conditions to determine a set of orthogonal functions, which generates a linear manifold of a Hilbert space, thus $f(x, t), \varphi_r(x)$ can be expressed in the Hilbert space.

References

- [1] G.Q. Bi, Applications of abstract operator in partial differential equation(i), *Pure and Applied Mathematics*. **13**(1997), 7-14.
- [2] G.Q. Bi, Applications of abstract operator in partial differential equation(ii), *Chinese Quarterly Journal of Mathematic*. **14**(1999), 80-87.
- [3] G.Q. Bi, Operator methods in high order partial differential equation, *Chinese Quarterly Journal of Mathematics*. **16**(2001), 88-101.
- [4] G.Q. Bi, Y.K. Bi, arXiv:1008.3808